

MDA’s MOVOR Data Hub provides insights into adoption of approved therapies for neuromuscular disease



Muscular Dystrophy Association

Elisabeth A Kilroy, PhD¹, Jessica Waits¹, Sharon Hesterlee, PhD¹

¹ Muscular Dystrophy Association, 161 N. Clark, Suite 3550, Chicago, IL 60601

Abstract

MOVOR as a Centralized Data Hub


The field of neuromuscular disease (NMD) research is experiencing tremendous growth as a result of progress in both diagnostics and therapeutics. However, there continues to be a critical data shortage, as data regarding disease progression have not been captured systematically across NMDs nor in centralized, accessible databases. The **neuroMuscular ObserVational Research Data Hub (MOVOR)** was created by the Muscular Dystrophy Association (MDA) to accelerate data collection, maximize drug development, and enhance the impact of new therapies. MOVOR is powered by the MDA Care Center network, which consists of over 150 multidisciplinary care centers across the United States. Currently, 60 care centers have activated the MOVOR Study Protocol and data from 3,948 participants and 12,376 encounters have been captured in MOVOR. Preliminary analyses of MOVOR Data have been performed to characterize the use FDA-approved therapies by participants living with spinal muscular atrophy (SMA), amyotrophic lateral sclerosis (ALS), or Duchenne muscular dystrophy (DMD). For participants living with SMA, the method of diagnosis (i.e., newborn screening, symptom onset) influences the type of therapy used and the age at which therapy is initiated. For participants living with ALS, over 75 percent of participants use an FDA-approved therapy, and these therapies are initiated on average less than 6 months after diagnosis. For participants living with DMD, the use of exon-skipping therapy is relatively limited among those participants who have mutations amenable to these therapies, with only 38 percent of participants receiving therapy. Future analyses will be performed to understand how these therapies may be impacting other core data elements captured by MOVOR, including medication usage, multidisciplinary referral types, and functional measures. As the first data hub to aggregate clinical and genetic data across multiple NMDs, **MOVOR continues to demonstrate its ability to transform the NMD space by serving as a powerful tool for clinical trial matching, clinical trial design and feasibility, and as a source for real-world data for pre- and post-approval submissions.**

About 10 years ago, MDA recognized that there was a significant data shortage in the NMD space and started crafting strategic approaches to accelerate data collection and its use by researchers, clinicians, and drug developers. One strategy that was identified was to leverage the MDA Care Center Network as a source for efficiently capturing clinical data and growing a longitudinal dataset. Each year, over 90,000 medical visits are conducted and over 60,000 individuals living with NMDs receive expert care at these centers.


60 Active MOVOR Sites



3,948 Participants



12,376 Encounters



Data elements captured by MOVOR are functional and disease-specific outcome measures that have been identified by KOLs as important to understanding disease mechanisms, tracking disease progression, and implementing standards of care.

31

core data elements captured by the electronic Case Report Forms (eCRFs)

Demographics eCRF [*] (During Enrollment)	Diagnosis eCRF [*] (During Enrollment)	Encounter eCRF [*] (During Clinical Visits)	Discontinuation eCRF [*] (After End of Study)
Disease Type Enrollment Date Gender DOB Race Ethnicity Insurance Education Employment	Age at Diagnosis Age at Symptom Onset Clinical Diagnosis First Symptoms Family History Genetic Testing Results	Encounter Date Height Weight Clinical Trial Participation Surgeries Hospitalizations Medications Pulmonary Devices Assistive Devices Functional Testing Pulmonary Tests Referral Types	Date of Discontinuation Reason for Discontinuation Date of Death Cause of Death


* Diagnosis and Encounter eCRFs contain additional unique fields for each indication.

Approved Therapy Use in DMD Participants


Approved Therapy Use in SMA Participants

Demographics of MOVOR Participants with DMD


99.6% of DMD participants are male




15.2 ± 6.7 years average age of active DMD participants



5.2 ± 3.6 years average age at diagnosis



93.1% of DMD participants have a genetic diagnosis



DMD Therapeutic Approval Timeline

There are 4 exon-skipping therapies approved by the FDA for individuals living with DMD.

2016

Exondys 51 (eteplirsen)
Sarepta
Mutations amenable to exon 51 skipping

2019

Vyondys 53 (golodirsen)
Sarepta
Mutations amenable to exon 53 skipping

2020


Viltepso (viltolarsen)
NS Pharma
Mutations amenable to exon 53 skipping

2021

Amondys 45 (casimersen)
Sarepta
Mutations amenable to exon 45 skipping

Therapy Use in MOVOR Participants with DMD

61.5% of DMD participants have a genetic deletion




38.3% of DMD participants harboring mutations amenable to exon-skipping are on therapy


Amenable Mutation	Amondys 45	Exondys 51	Vyondys 53	Viltepso
No Therapy	32 (84%)	13 (23%)	23 (77%)	27 (90%)
Therapy	6 (16%)	38 (68%)	7 (23%)	3 (10%)
No Mutation Information	0	5 (9%)	0	0

Demographics of MOVOR Participants with SMA


53.7% of SMA participants are male




17.9 ± 17.3 years average age of active SMA participants



64.2% of SMA participants were diagnosed after symptom onset



92.2% of SMA participants have a genetic confirmation



SMA Therapeutic Approval Timeline

There are 3 therapies approved by the FDA for individuals living with SMA.

2016

Spinraza (nusinersen)
Biogen
Antisense oligonucleotide that selectively binds and targets RNA to regulate SMN protein production

2019


Zolgensma (onasemnogene abeparvovec-xioi)
Novartis
Gene therapy that introduces a fully functional copy of the human SMN gene

2020


Evrysdi (risdiplam)
Genentech
SMN2-splicing modifier that increases the production of full-length SMN protein

Therapy Use in MOVOR Participants with SMA


69.7% of SMA participants have used an FDA-approved therapy




57.0% of SMA participants begin therapy before the age of 10



60.1% of SMA participants have received Spinraza



SMA participants diagnosed via prenatal screening start therapy sooner




Therapy Start Age (Years)				Therapy Start Age (Years)					
	Zolgensma	Evrysdi	Spinraza		Clinical Presentation	Sibling Diagnosis	Newborn Screening	Prenatal Screening	Unknown
count	43.00	46.00	134.00	count	140	10	51	10	12
mean	0.58	15.93	15.77	mean	17.24	10.00	1.49	0.40	23.08
std	0.73	14.13	17.10	std	15.87	22.57	3.80	0.70	17.65


Approved Therapy Use in ALS Participants

Demographics of MOVOR Participants with ALS


56.9% of ALS participants are male




64.5 ± 10.9 years average age of active ALS participants



1.5 ± 1.3 years average time from symptom onset to diagnosis



72.5% of ALS participants had non-Bulbar onset



ALS Therapeutic Approval Timeline

There are 5 therapies approved by the FDA for individuals living with ALS.

1995

Rilutek (riluzole)
Sanofi
Inhibits glutamate release

2011

Nuedexta
Avanir Pharmaceuticals
Treatment of pseudobulbar affect

2017

Radicava (edaravone)
Mitsubishi Tanabe Pharma America
Prevent accumulation of reactive oxygen species

2018


Tiglutik (thickened riluzole)
ITF Pharma
Inhibits glutamate release

2019


Exservan (riluzole oral film)
Mitsubishi Tanabe Pharma America
Inhibits glutamate release

Therapy Use in MOVOR Participants with ALS


77.8% of ALS participants have used an FDA-approved therapy



5.2 ± 15.6 months average time from diagnosis to start an FDA-approved therapy



Regional Differences in FDA-approved therapy use are not present



Conclusions

The MOVOR Data Hub is experiencing tremendous growth in the number of participants enrolled in this observational study as well as in the number of longitudinal clinical encounters. Standardization of data elements across the 7 indications provides MOVOR with the ability to examine the same outcome measures across time. For these analyses, we examined the use of FDA-approved therapies in MOVOR Participants living with DMD, SMA, and ALS. The majority of ALS and SMA participants are receiving (or have received) an FDA-approved therapy but relatively few DMD participants have received a therapy. Building these two cohorts (therapy vs therapy-naïve) provides an opportunity to explore how therapies impact disease progression as well as standard of care practices. The MOVOR Data Hub is a valuable tool for monitoring longitudinal response to therapies and could serve as an excellent source for real-world efficacy data for future regulatory submissions.

Contact the MOVOR Team



Sharon Hesterlee, PhD
EVP & Chief Research Officer
shesterlee@mداusa.org



Elisabeth Kilroy, PhD
MOVOR Data Analytics
ekilroy@mداusa.org



Jessica Waits
MOVOR Clinical Operations
jwaits@mداusa.org

For access to MOVOR Data, please email MDAMOVOR@mداusa.org

Acknowledgements
This study was conducted using data from the Muscular Dystrophy Association’s neuroMuscular ObserVational Research (MOVOR) Data Hub. MOVOR is operated through participating MDA Care Centers with the support of participants, MOVOR Site Pls, coordinators, and staff. The authors would also like to thank members of the MOVOR Research Advisory Committee, including James Berry, Barry Byrne, John Day, Rod Howell, Monkol Lek, Nicholas Maragakis, and Sabrina Paganoni, for sharing their expertise in neuromuscular diseases, the original MDA team members who spear-headed the development of the MOVOR eCRFs, including Grace Pavlath, Lianna Orlando, Amanda Hayden-Phillips, and Laura Hagerty, and Tracy Blumenfeld for her contributions in launching MOVOR. Finally, the authors would like to thank the teams at IQVIA and DNAexus.